

HOMOTOPIES OF LIE CROSSED MODULE MORPHISMS

İLKER AKÇA AND YAVUZ SIDAL

ABSTRACT. In this paper we will define notion homotopy of morphisms of crossed modules of Lie algebras. Then we construct a groupoid structure of Lie crossed module morphisms and their homotopies.

1. INTRODUCTION

Crossed modules were firstly introduced by J.H.C Whitehead in his work on combinatorial homotopy theory [6]. They have found important role in many areas of mathematics (including homotopy theory, homology and cohomology of groups, algebraic K-theory, cyclic homology, combinatorial group theory and differential geometry). Kassel and Loday [4] introduced crossed modules of Lie algebras as computational algebraic objects equivalent to simplicial Lie algebras with associated Moore complex of length 1.

The homotopy relation between crossed module morphisms $\mathcal{P} \rightarrow \mathcal{P}'$ can be equivalently addressed either by considering natural functorial path objects for $\mathcal{P}'$ or cylinder objects for $\mathcal{P}$. It yields, given any two crossed modules $\mathcal{P}$ and $\mathcal{P}'$, a groupoid of morphisms $\mathcal{P} \rightarrow \mathcal{P}'$ and their homotopies. In addition, the homotopy relation between crossed module morphisms $\mathcal{P} \rightarrow \mathcal{P}'$ is an equivalence relation in the general case, with no restriction on $\mathcal{P}$ or $\mathcal{P}'$. This should be compared with what would be guaranteed from the model category [3] point of view, where we would expect homotopy of maps $\mathcal{P} \rightarrow \mathcal{P}'$ to be an equivalence relation only when $\mathcal{P}$ is a cofibrant (given that any object is fibrant). In the well-known model category structure in the category of crossed modules [2], obtained by transporting the usual model category structure of the category of simplicial sets, $\mathcal{P} = (\partial : M \rightarrow P)$ is cofibrant if and only if P is a free group ([5]).

Whitehead in [6] explored homotopies of morphisms of his "homotopy systems" and this was put in the general context of crossed complexes of groupoids by Brown and Higgins in [1]. In this paper we will define notion of homotopy for morphisms of crossed modules of Lie algebras and we will show that if $\mathcal{P}$ and $\mathcal{P}'$ are crossed modules of Lie algebras, without any restriction on $\mathcal{P}$ or $\mathcal{P}'$, then we have a groupoid of crossed module morphisms $\mathcal{P} \rightarrow \mathcal{P}'$ and their homotopies, similarly to the group case.

2. CROSSED MODULES

J.H.C Whitehead (1949) [6] described crossed modules in various contexts especially in his investigations into the algebraic structure of relative homotopy groups.

2010 *Mathematics Subject Classification.* 18D05; 18G55; 18D20.
Key words and phrases. Crossed Modules, derivations, homotopy.

In this section, we recall the definition of crossed modules of Lie algebras given by Kassel and Loday [4].

Let M and P be two Lie algebras. By an action of P on M we mean a bilinear map $P \times M \longrightarrow M$, $(p, m) \longmapsto p \cdot m$ satisfying

$$[p, p'] \cdot m = p \cdot (p' \cdot m) - p' \cdot (p \cdot m)$$

$$p \cdot [m, m'] = [p \cdot m, m'] + [m, p \cdot m']$$

for all $m, m' \in M$, $p, p' \in P$. For instance, if P is a subalgebra of some Lie algebra Q (including possibly the case $P = Q$), and if M is an ideal in Q , then Lie bracket in Q yields an action of P on M .

A crossed module of Lie algebras is a Lie homomorphism $\partial : M \longrightarrow P$ together with an action of P on M such that for all $m, m' \in M$, $p \in P$

$$\mathbf{CM1)} \quad \partial(p \cdot m) = [p, \partial(m)]$$

and

$$\mathbf{CM2)} \quad \partial m \cdot m' = [m, m'] .$$

The last condition is called the Peiffer identity. We denote such a crossed module by $\mathcal{P} = (M, P, \partial)$.

A morphism of crossed modules of Lie algebras from $\mathcal{P} = (M, P, \partial)$ to $\mathcal{P}' = (M', P', \partial')$ is a pair of Lie algebra morphisms,

$$\theta : M \longrightarrow M', \psi : P \longrightarrow P'$$

such that

$$\theta(p \cdot m) = \psi(p) \cdot \theta(m) \text{ and } \partial' \theta(m) = \psi \partial(m).$$

Therefore we can define the category of Lie crossed modules denoting it as **LXmod**.

2.1. Examples. 1. Let I be any ideal of a Lie algebra P . Consider an inclusion map

$$inc : I \longrightarrow P.$$

Then (I, P, inc) is a crossed module. Conversely given any crossed module $\partial : M \longrightarrow P$, one can easily verify that $\partial M = I$ is an ideal in P .

2. Let M be any R -bimodule. It can be considered as an R -algebra with zero multiplication, and then $\mathbf{0} : M \longrightarrow R$ is a crossed R -module by $\mathbf{0}(c) \cdot c' = \mathbf{0}c' = 0 = cc'$ for all $c, c' \in M$.

Conversely, given any crossed module $\partial : C \longrightarrow R$, then $Ker \partial$ is an $R/\partial C$ -module.

3. HOMOTOPIES OF LIE CROSSED MODULE MORPHISMS

Whitehead in [6] explored homotopies of morphisms of his "homotopy systems" and this was put in the general context of crossed complexes of groupoids by Brown and Higgins in [1]. In this section we define notion of homotopy for morphisms of crossed modules of Lie algebras.

Definition 3.1. Let $\mathcal{P} = (M, P, \partial)$ and $\mathcal{P}' = (M', P', \partial')$ be crossed modules of Lie algebras, $f = (f_1, f_0)$ and $g = (g_1, g_0)$ be crossed module morphisms $\mathcal{P} \rightarrow \mathcal{P}'$. If there is a linear map $d : P \rightarrow M'$ such that,

$$\begin{aligned} g_0(p) &= f_0(p) + \partial' d(p) \\ g_1(m) &= f_1(m) + d\partial(m) \end{aligned}$$

for $m \in M$ and $p \in P$, then we say that d is a homotopy connecting f to g and we write $d : f \simeq g$ or $f \xrightarrow{d} g$.

Definition 3.2. Let $\mathcal{P} = (M, P, \partial)$ and $\mathcal{P}' = (M', P', \partial')$ be crossed modules of Lie algebras and $f = (f_1, f_0)$ be a crossed module morphism $\mathcal{P} = (M, P, \partial) \rightarrow \mathcal{P}' = (M', P', \partial')$. Then a linear map $d : P \rightarrow M'$ satisfying for all $p, p' \in P$

$$d[p, p'] = f_0(p) \cdot d(p') - f_0(p') \cdot d(p) + [d(p), d(p')]$$

is called an f_0 -derivation .

Proposition 3.3. Let $\mathcal{P} = (M, P, \partial)$ and $\mathcal{P}' = (M', P', \partial')$ be crossed modules of Lie algebras and $f = (f_1, f_0)$ be a crossed module morphism $\mathcal{P} \rightarrow \mathcal{P}'$. If $d : P \rightarrow M'$ is an f_0 -derivation, then the maps $g_0 : P \rightarrow P'$ defined by $g_0(p) = f_0(p) + \partial' d(p)$ for all $p \in P$ and $g_1 : M \rightarrow M'$ defined by $g_1(m) = f_1(m) + d\partial(m)$ for all $m \in M$ are Lie algebra morphisms.

Proof. For $p, p' \in P$,

$$\begin{aligned} g_0(p + p') &= f_0(p + p') + \partial' d(p + p') \\ &= f_0(p) + f_0(p') + \partial'(d(p) + d(p')) \\ &= f_0(p) + f_0(p') + \partial' d(p) + \partial' d(p') \\ &= g_0(p) + g_0(p'), \end{aligned}$$

and

$$\begin{aligned} g_0[p, p'] &= f_0[p, p'] + \partial' d[p, p'] \\ &= [f_0(p), f_0(p')] + \partial'(f_0(p) \cdot d(p') - f_0(p') \cdot d(p) \\ &\quad + [d(p), d(p')]) \\ &= [f_0(p), f_0(p')] + \partial'(f_0(p) \cdot d(p')) - \partial'(f_0(p') \cdot d(p)) \\ &\quad + \partial'[d(p), d(p')] \\ &= [f_0(p), f_0(p')] + [f_0(p), \partial' d(p')] - [f_0(p'), \partial' d(p)] \\ &\quad + [\partial' d(p), \partial' d(p')] \\ &= [f_0(p), f_0(p')] + [f_0(p), \partial' d(p')] + [\partial' d(p), f_0(p'),] \\ &\quad + [\partial' d(p), \partial' d(p')] \\ &= [f_0(p) + \partial' d(p), f_0(p') + \partial' d(p')], \\ &= [g_0(p), g_0(p')]. \end{aligned}$$

Thus g_0 is a Lie algebra morphism.

For $m, m' \in M$,

$$\begin{aligned} g_1(m + m') &= f_1(m + m') + d\partial(m + m') \\ &= f_1(m) + f_1(m') + d(\partial(m) + \partial(m')) \\ &= f_1(m) + f_1(m') + d\partial(m) + d\partial(m') \\ &= g_1(m) + g_1(m'), \end{aligned}$$

and

$$\begin{aligned}
g_1[m, m'] &= f_1[m, m'] + d\partial[m, m'] \\
&= [f_1(m), f_1(m')] + d[\partial(m), \partial(m')] \\
&= [f_1(m), f_1(m')] + f_0(\partial(m)) \cdot d(\partial(m')) \\
&\quad - f_0(\partial(m')) \cdot d(\partial(m)) + [d(\partial(m)), d(\partial(m'))] \\
&= [f_1(m), f_1(m')] + \partial' f_1(m) \cdot d(\partial(m')) \\
&\quad - \partial' f_1(m') \cdot d(\partial(m)) + [d(\partial(m)), d(\partial(m'))] \\
&= [f_1(m), f_1(m')] + [f_1(m), d(\partial(m'))] - [f_1(m'), d(\partial(m))] \\
&\quad + [d(\partial(m)), d(\partial(m'))] \\
&= [f_1(m), f_1(m')] + [f_1(m), d(\partial(m'))] + [d(\partial(m)), f_1(m')] \\
&\quad + [d(\partial(m)), d(\partial(m'))] + [d(\partial(m)), d(\partial(m'))] \\
&= [f_1(m) + d(\partial(m)), f_1(m') + d(\partial(m'))] \\
&= [g_1(m), g_1(m')]
\end{aligned}$$

thus g_1 is a Lie algebra morphism. $\square$

Proposition 3.4.

$$g_0\partial = \partial'g_1.$$

Proof. For $m \in M$,

$$\begin{aligned}
(g_0\partial)(m) &= g_0(\partial(m)) \\
&= f_0(\partial(m)) + \partial'd(\partial(m)) \\
&= \partial'(f_1(m)) + \partial'(d\partial(m)) \\
&= \partial'(f_1(m) + d\partial(m)) \\
&= \partial'(g_1(m)) \\
&= (\partial'g_1)(m),
\end{aligned}$$

so

$$g_0\partial = \partial'g_1. \quad \square$$

Proposition 3.5. For $p \in P$ and $m \in M$,

$$g_1(p \cdot m) = g_0(p) \cdot g_1(m).$$

Proof.

$$\begin{aligned}
g_1(p \cdot m) &= f_1(p \cdot m) + d\partial(p \cdot m) \\
&= f_0(p) \cdot f_1(m) + d[p, \partial(m)] \\
&= f_0(p) \cdot f_1(m) + f_0(p) \cdot d\partial(m) - f_0\partial(m) \cdot d(p) + [d(p), d\partial(m)] \\
&= f_0(p) \cdot f_1(m) + f_0(p) \cdot d\partial(m) - \partial' f_1(m) \cdot d(p) + [d(p), d\partial(m)] \\
&= f_0(p) \cdot f_1(m) + f_0(p) \cdot d\partial(m) - [f_1(m), d(p)] + [d(p), d\partial(m)] \\
&= f_0(p) \cdot f_1(m) + f_0(p) \cdot d\partial(m) + [d(p), f_1(m)] + [d(p), d\partial(m)] \\
&= f_0(p) \cdot f_1(m) + f_0(p) \cdot d\partial(m) + [d(p), f_1(m) + d\partial(m)] \\
&= f_0(p) \cdot f_1(m) + f_0(p) \cdot d\partial(m) + \partial'd(p) \cdot (f_1(m) + d\partial(m)) \\
&= g_0(p) \cdot g_1(m).
\end{aligned}$$

$\square$

Thus by the above propositions, we can give the following theorem.

Theorem 3.6. *Let $\mathcal{P} = (M, P, \partial)$ and $\mathcal{P}' = (M', P', \partial')$ be crossed modules of Lie algebras and $f = (f_1, f_0)$ be a crossed module morphism $\mathcal{P} \rightarrow \mathcal{P}'$. If $d : P \rightarrow M'$ is an f_0 -derivation, then the map $g = (g_1, g_0) : \mathcal{P} \rightarrow \mathcal{P}'$ is a Lie crossed module morphism.*

Corollary 3.7. *Let $\mathcal{P} = (M, P, \partial)$ and $\mathcal{P}' = (M', P', \partial')$ be crossed modules of Lie algebras and $f = (f_1, f_0), g = (g_1, g_0)$ be crossed module morphisms $\mathcal{P} \rightarrow \mathcal{P}'$. Then the f_0 -derivation $d : P \rightarrow M'$ satisfying for all $p, p' \in P$*

$$d[p, p'] = f_0(p) \cdot d(p') - f_0(p') \cdot d(p) + [d(p), d(p')]$$

is a homotopy connecting f to g .

4. GROUPOID STRUCTURE FOR CROSSED MODULE MORPHISMS AND THEIR HOMOTOPIES

In this section we construct a groupoid structure whose objects are the crossed module morphisms $\mathcal{P} \rightarrow \mathcal{P}'$, with morphisms being the homotopies between them.

Lemma 4.1. *Let $\mathcal{P} = (M, P, \partial)$ and $\mathcal{P}' = (M', P', \partial')$ be crossed modules of Lie algebras and $f = (f_1, f_0)$ be a crossed module morphism $\mathcal{P} \rightarrow \mathcal{P}'$. Then the null function $0 : P \rightarrow M'$, $0(p) = 0_{M'}$ defines an f_0 derivation connecting f to f .*

Lemma 4.2. *Let $f = (f_1, f_0)$ and $g = (g_1, g_0)$ be crossed module morphisms $\mathcal{P} \rightarrow \mathcal{P}'$ and d be an f_0 -derivation connecting f to g . Then the linear map $\bar{d} = -d : P \rightarrow M'$ with $\bar{d}(p) = -d(p)$ is a g_0 derivation connecting g to f .*

Proof. Since d is an f_0 derivation connecting f to g , we have

$$(4.1) \quad f_0(p) = g_0(p) + \partial' \bar{d}(p) \text{ and } f_1(m) = g_1(m) + \bar{d} \partial(m).$$

Moreover $\bar{d}$ is a g_0 derivation, since:

$$\begin{aligned} \bar{d}[p, p'] &= -d[p, p'] \\ &= -(f_0(p) \cdot d(p') - f_0(p') \cdot d(p) + [d(p), d(p')]) \\ &= -f_0(p) \cdot d(p') + f_0(p') \cdot d(p) - [d(p), d(p')] \\ &\quad - [d(p), d(p')] + [d(p), d(p')] \\ &= -f_0(p) \cdot d(p') - \partial' d(p) \cdot d(p') + f_0(p') \cdot d(p) \\ &\quad + \partial' d(p') \cdot d(p) + [d(p), d(p')] \\ &= f_0(p) \cdot (-d(p')) + \partial' d(p) \cdot (-d(p')) - f_0(p') \cdot (-d(p)) \\ &\quad - \partial' d(p') \cdot (-d(p)) + [-d(p), -d(p')] \\ &= (f_0(p) + \partial' d(p)) \cdot (-d(p')) - (f_0(p') + \partial' d(p')) \cdot (-d(p)) \\ &\quad + [-d(p), -d(p')] \\ &= g_0(p) \cdot (-d(p')) - g_0(p') \cdot (-d(p)) + [-d(p), -d(p')]. \end{aligned}$$

□

Lemma 4.3. *(Concatenation of derivations) Let f, g and h be Lie crossed module morphisms $\mathcal{P} \rightarrow \mathcal{P}'$, d be an f_0 derivation connecting f to g , and d' be a g_0 derivation connecting g to h . Then the linear map $(d + d') : P \rightarrow M'$ such that $(d + d')(p) = d(p) + d'(p)$, defines an f_0 derivation (therefore a homotopy) connecting f to h .*

Proof. We know that $f \xrightarrow{(f_0, d)} g$ and $g \xrightarrow{(g_0, d')} h$. Therefore by definition

$$\begin{aligned} h_0(p) &= f_0(p) + \partial'(d + d')(p) \\ h_1(m) &= f_1(m) + (d + d')\partial(m). \end{aligned}$$

Let us see that $d + d'$ satisfies the condition for it to be an f_0 derivation:

$$\begin{aligned}
(d + d')[p, p'] &= d[p, p'] + d'[p, p'] \\
&= f_0(p) \cdot d(p') - f_0(p') \cdot d(p) + [d(p), d(p')] \\
&\quad + g_0(p) \cdot d'(p') - g_0(p') \cdot d'(p) + [d'(p), d'(p')] \\
&= f_0(p) \cdot d(p') - f_0(p') \cdot d(p) + [d(p), d(p')] \\
&\quad + (f_0(p) + \partial' d(p)) \cdot d'(p') - (f_0(p') + \partial' d(p')) \cdot d'(p) \\
&\quad + [d'(p), d'(p')] \\
&= f_0(p) \cdot d(p') - f_0(p') \cdot d(p) + [d(p), d(p')] + f_0(p) \cdot d'(p') \\
&\quad + \partial' d(p) \cdot d'(p') - f_0(p') \cdot d'(p) - \partial' d(p') \cdot d'(p) \\
&\quad + [d'(p), d'(p')] \\
&= f_0(p) \cdot d(p') - f_0(p') \cdot d(p) + [d(p), d(p')] + f_0(p) \cdot d'(p') \\
&\quad + [d(p), d'(p')] - f_0(p') \cdot d'(p) - [d(p'), d'(p)] + [d'(p), d'(p')] \\
&= f_0(p) \cdot (d(p') + d'(p')) - f_0(p') \cdot (d(p) + d'(p)) \\
&\quad + [d(p) + d'(p), d(p') + d'(p')]
\end{aligned}$$

for all $p, p' \in P$. Therefore $(d + d')$ is an f_0 derivation connecting f to h . $\square$

Theorem 4.4. *Let $\mathcal{P}$ and $\mathcal{P}'$ be two arbitrary crossed modules of Lie algebras. We have a groupoid $HOM(\mathcal{P}, \mathcal{P}')$, whose objects are the crossed module morphisms $\mathcal{P} \rightarrow \mathcal{P}'$, the morphisms being their homotopies. In particular the relation below, for crossed module morphisms $\mathcal{P} \rightarrow \mathcal{P}'$, is an equivalence relation:*

$$f \simeq g \Leftrightarrow \text{there exists an } f_0 \text{ derivation } d \text{ connecting } f \text{ to } g.$$

5. ACKNOWLEDGEMENTS

This research was supported by Eskişehir Osmangazi University Scientific Research Center (BAP) under Grant No:2016-1129.

REFERENCES

- [1] BROWN, R. AND HIGGINS P. J., Tensor Products and Homotopies for ω -groupoids and crossed complexes, *Journal of Pure and Applied Algebra* **47**, (1987), 1-33.
- [2] CABELLO, J.G. AND GARZON A.R. Closed model structures for algebraic models of n-types, *Journal of Pure and Applied Algebra* 103 (**3**), (1995), 287-302.
- [3] DWYER, W.G. - SPALINSKI, J. Homotopy theories and model categories, In *Handbook of algebraic topology*, pages 73-126. Amsterdam: Nort Holland, (1995).
- [4] KASEL, C. and LODAY, J.L. Extensions centrales d'algebres de Lie. *Ann. Inst. Fourier (Grenoble)*, **33**, (1982) 119-142.
- [5] NOOHI, B. Notes on 2-groupoids, 2-groups and crossed modules, *Homology Homotopy Appl.* 9 (1), (2007), 75-106.
- [6] WHITEHEAD, J.H.C. Combinatorial Homotopy I and II, *Bull. Amer. Math. Soc.*, **55**, 231-245 and 453-456 (1949).

Current address: Department of Mathematics and Computer Science, Art and Science Faculty, Eskişehir Osmangazi University, 26480, Eskişehir, Turkey
E-mail address: iakca@ogu.edu.tr

Current address: Isiklar Air Force High School , Maths Teacher, 16039, Bursa/TURKEY
E-mail address: yavuz.sidal@yahoo.com